

Evidence for a conformal phase in $SU(N)$ gauge theories

A. Deuzeman,¹ M.P. Lombardo,² and E. Pallante¹

¹*Centre for Theoretical Physics, University of Groningen, 9747 AG, Netherlands*

²*INFN-Laboratori Nazionali di Frascati, I-00044, Frascati (RM), Italy*

(Dated: December 21, 2009)

We discuss the existence of a conformal phase in $SU(N)$ gauge theories in four dimensions. Simulations are carried out with three colors and twelve flavors of dynamical staggered fermions in the fundamental representation. The analysis of the chiral order parameter and the mass spectrum of the theory indicates the restoration of chiral symmetry at zero temperature and the presence of a Coulomb phase, depicting a scenario compatible with the existence of an infrared stable fixed point at nonzero coupling. Our analysis thus supports the conclusion that the onset of the conformal window for QCD-like theories is smaller than $N_f = 12$, before the loss of asymptotic freedom at sixteen and a half flavors.

With the imminent activity of the LHC experiments and the quest for a theory describing fundamental forces beyond the electroweak symmetry breaking scale, renewed interest has arisen in the most elusive aspects of gauge theories. In particular, the possibility of an emergent quasi-conformal symmetry in theories with fermionic content has attained a strong experimental appeal. There are multiple reasons for pursuing this search. Resolving conformal behavior would complete our understanding of the phase diagram of gauge theories by varying temperature and number of flavors, as sketched in Fig. 1. It sheds light on how the low temperature and large flavor number quasi-conformal phase may be connected to the high temperature and low flavor number quark-gluon plasma phase. It is essential for theoretically establishing or excluding walking technicolor-type theories and more generally strongly interacting dynamics above the electroweak symmetry breaking scale. Finally, elucidating the way conformal symmetry or its remnants drive the dynamics of particle interactions with or without supersymmetry contributes to clarifying the possible connection of field theory to string theory that the AdS/CFT correspondence seems to imply.

In the early eighties our understanding of the perturbative behavior of non Abelian gauge theories was enriched by two seminal papers [1, 2]. It was noticed that a second zero of the two-loop beta function of an $SU(3)$ gauge theory with N_f massless fermions in the fundamental representation appears for $N_f \gtrsim 8.05$, at $g^{*2} \neq 0$, before the loss of asymptotic freedom at $N_f^c = 16\frac{1}{2}$. This fact implies, at least perturbatively, the appearance of an infrared fixed point (IRFP). The fixed point moves closer to zero coupling as the number of flavors approaches N_f^c . The dynamics of chiral symmetry have led to the discovery of the conformal window in QCD-like theories [3, 4]: chiral symmetry breaking can only occur below a critical number of flavors N_f^* . Between N_f^* and N_f^c the conformal window opens up. Finding the actual value of the critical number of flavors N_f^* at which chiral symmetry breaking takes place is a non-perturbative problem for which the lattice approach is the method of choice. Re-




FIG. 1: A projected view of the phase diagram of QCD-like theories in the temperature (T), flavor number (N_f) and bare coupling (g) space. In the T - N_f plane, the critical line is a phase boundary between the chirally broken hadronic phase and the chirally symmetric quark gluon plasma, the zero temperature end point of which is the onset of the conformal window. The zero temperature projected plane is inspired by the scenario in Refs. [3, 4], see Fig. 2.

cent studies have provided evidence that $N_f = 8$ lies within the hadronic phase of QCD [5, 6, 7, 8, 9]. A recent study of the $SU(3)$ running coupling [5, 8] by use of the lattice Schrödinger functional has concluded that $N_f = 12$ should already be in the conformal window. Other numerical studies, however, challenged this conclusion [10, 11, 12]. This is hardly surprising, given that $N_f = 12$ should be very close to the critical number of flavors [13, 14, 15, 16], making a numerical study particularly delicate.

The current strategy, complementary to that of Refs. [5, 8], is inspired by the physics of phase transitions; it allows for the exploration of multiple aspects of the theory in different regimes and regions of the phase diagram, in order to probe the existence and properties of an IRFP inside and outside its basin of attraction.

A lattice strategy The strategy of this study has received heuristic guidance by the scenario depicted in




FIG. 2: (color online) Phase diagram of an $SU(3)$ gauge theory with fundamental fermions in the N_f - g plane after Ref. [4]. Theories for $N_f < N_f^*$ are QCD-like in the continuum, while for $N_f^* < N_f < N_f^c$ develop a conformal phase. S and A refer to chirally symmetric and asymmetric, respectively. The dashed (green) line qualitatively indicates the location of the Banks-Zaks IRFP [2]. The dot-dashed (red) line indicates a lattice bulk transition, which has been observed at $N_f = 12$ and $N_f = 16$. It stops at some small N_f value. The line at $N_f = N_f^*$ represents the conformal phase transition [3, 4], which is absent in the original Banks-Zaks scenario. The beta function on the conformal side is also sketched.

Refs. [3, 4] and sketched in Fig. 2. We will implicitly assume the validity of that scenario. Following Fig. 2, at a given $N_f > N_f^*$ and increasing the coupling from $g = 0$, one crosses the conformal line, location of the IRFP's, going from a chirally symmetric (S) and asymptotically free phase (quasi-conformal phase) to a symmetric, but not asymptotically free one (Coulomb or QED-like phase). A phase transition need not be associated with the line of IRFP's, differently from what was originally speculated in Ref. [2]. At even larger couplings, a transition to a strongly coupled chirally asymmetric (A) phase will always occur in the lattice regularized theory. The latter is referred to as a bulk phase transition. In the symmetric phases at nonzero coupling the conformal symmetry is still broken by ordinary perturbative contributions. They generate the running of the coupling constant which is different on the two sides of the symmetric phase. See Ref. [4] for a detailed discussion of this point.

A theory in the hadronic phase, $N_f < N_f^*$, has a thermal phase transition in the continuum from a low temperature chirally broken phase to a high temperature chirally symmetric – quark gluon plasma – phase. Thus, as argued in Ref. [6], the observation of a thermal transition in the continuum limit is incompatible with the existence of a conformal fixed point, see Fig. 1. It is also clear from Fig. 2 that the presence of a Coulomb phase next to the bulk transition at weaker coupling is also a distinguishing feature of the conformal window. Here, the non-perturbative beta function should be positive, implying a weakening of the effective coupling over increasing distances. The appearance of such a region is,




FIG. 3: (color online) The bulk transition in the chiral condensate for $am = 0.05$ on lattices of $16^3 \times 8$ (circles), and 16^4 (crosses) as a function of the bare lattice coupling g_L . Data are shown in the range $6/g_L^2 = 2.5$ to 4.7 . The location of the transition is identical, while the curves describe physics at temperatures differing by a factor of two. Simulation errors are within symbol size.

in principle, a sufficient condition for the existence of an IRFP, since the perturbative beta function of $SU(3)$ with $N_f < 16\frac{1}{2}$ in the extreme weak coupling regime is known to be negative. Note, however, that the beta function is not universal away from fixed points with diverging correlation lengths and one can therefore not exclude *a priori* the appearance of spurious fixed points at intermediate values of the coupling constant [17].

The evidence presented here thus consists of a few components. First, it will be demonstrated that the location of the transition from the chirally symmetric to the broken phase is not sensitive to the physical temperature and is therefore compatible with a bulk nature. Subsequently, we will present a detailed study of the mass dependence of the chiral condensate on the weak coupling side of the bulk transition, which clearly favors exact chiral symmetry. Finally, the behavior of the mass spectrum close to the bulk transition will be studied, and found to be compatible with a positive beta function, similarly to the observations of Ref. [17] for $N_f = 16$. These results are consistent with the scenario for conformality of Fig. 2.

Results We performed simulations of an $SU(3)$ gauge theory with twelve flavors of staggered fermions in the fundamental representation. The simulations were done using a tree level Symanzik improved gauge action to suppress lattice artifacts, and Kogut-Susskind (staggered) fermions with the Naik improvement scheme, that effectively extends the Symanzik improvement to the matter content. The order parameter of the chiral phase transition for a QCD-like theory is the chiral condensate $\langle\bar{\psi}\psi\rangle$. We underscore that staggered fermions have a remnant of exact chiral symmetry which allows a precise definition of the order parameter also on a coarse lattice. Fig. 3 displays a sudden variation of the chiral order parameter as a function of the bare lattice coupling constant g_L , and for two values of the temporal lattice extent N_t differing




FIG. 4: (color online) Deviations of the predicted value for an extrapolation to the chiral limit of the chiral condensate for $6/g_L^2 = 3.9$ (circles) and $6/g_L^2 = 4.0$ (triangles), fitting to a linear form $Am + B$ (red) - with m the bare mass in lattice units - and a power law Am^δ (blue), rescaled by the standard deviation of each measurement. The data and corresponding fits are displayed in the inset on linear scales.

by a factor of two. The temperature of the system is related to the lattice temporal extent as $T = 1/a(g_L)N_t$, with $a(g_L)$ the lattice spacing for a given lattice coupling. From Fig. 3 one infers that the transition – or rapid crossover – happens at identical values of the critical coupling $g_L^c = 1.35(3)$, thus implying they occur at vastly different physical temperatures. One concludes that the observed phase transition is driven purely by the bare coupling constant itself and is therefore of bulk nature. Further information on this phase transition might shed light on the occurrence of a conjectured ultraviolet fixed point at strong coupling in the continuum theory [18].

To discriminate between a weakly broken and an exact chiral symmetry on the weak coupling side of the bulk transition, we performed a high statistics study of the mass dependence of the chiral condensate for two selected couplings. Measurements were performed on three different volumes for each mass, up to 32^4 , and the difference between the largest two volumes found to be smaller than both the difference between the smallest volumes and the statistical uncertainty in all measurements. Data displayed in Fig. 4, all at volume 24^4 , can be considered as infinite volume measurements within their errors. We fitted the chiral condensate data to $\langle \bar{\psi}\psi \rangle = Am + \langle \bar{\psi}\psi \rangle_0$, appropriate for a chirally broken ($\langle \bar{\psi}\psi \rangle_0 \neq 0$) or symmetric phase, away from criticality and in the absence of an IRFP, and to a power law $\langle \bar{\psi}\psi \rangle = Am^\delta$. The latter should be expected in a symmetric phase and in the vicinity of a fixed point with anomalous dimensions. For instance, this behavior was observed for lattice QED in Ref. [19]. The pattern of the deviations in the linear fits indicates a significant curvature, which is reflected in the quality of the fit. The χ^2 per d.o.f. was 0.57 (power) as compared to 3.21 (linear) for $g_L = 1.240$ ($6/g_L^2 = 4.0$), and 0.86 as compared to

3.81 for $g_L = 1.224$ ($6/g_L^2 = 3.9$), with best fitted exponent $\delta = 0.965(1)$ for both. A power law fit with a free intercept always produced a best fitted value consistent with zero. To complete our study of the symmetric phase, following Ref. [17], we measured the mass spectrum and used the results to determine the lines of constant physics in the two dimensional parameter space g_L and am , the bare quark mass of degenerate fermions. Along these lines the coupling and masses are all functions of the lattice spacing a . Since all dimensionful quantities measured on the lattice will be expressed in terms of the lattice spacing and will therefore vary with g_L even if they do not physically, a dimensionless quantity has to be taken as a reference. A convenient choice is the ratio of the ground state excitations in the pseudoscalar and vector channels, with slight abuse of nomenclature from QCD referred to as the π and ρ masses. In Fig. 5(a) we report on the measured values of m_π and m_ρ as a function of the bare coupling and the bare mass. Simulations were done on $16^3 \times 24$ volumes, while a set of measurements at larger volumes showed that finite volume effects were under control.

In Fig. 5(b) we show the ratios of measured pseudoscalar and vector masses, for a fixed coupling and as a function of the bare quark mass. We have superimposed the ratios of the best fits to the raw mass data, confirming the good quality of the interpolations derived in Fig. 5(a) and later used to produce Fig. 6. It is immediately evident that, in order to keep the ratio constant, we should simultaneously decrease the lattice coupling g_L and increase the bare mass. These results already indicate that the lattice spacing increases while decreasing the coupling. This is the same behavior as observed for $N_f = 16$, and of the pion to sigma ratio in QED. It is also expected of a one parameter beta function with a positive sign. To refine the analysis, and express the result in terms of a physical observable, we proceed as follows. Given the ratio m_π/m_ρ at reference values of g_L and am , one can determine a value $a'm'$ at coupling g'_L in the surroundings of g_L that reproduces the same ratio and thus lies on the same line of constant physics. This is implemented by fitting the measured values of both masses to a parametrization, as shown in Fig. 5(a), then determining the isolines from the ratio of the two parametrizations. The physical pseudoscalar mass being constant along each of these lines, the ratio of measured values $am_\pi/a'm_\pi$ directly determines the ratio of the lattice spacings a/a' . If a decrease in g_L is associated with an increase in the lattice spacing, the sign of the beta function is positive, i.e. that of the Coulomb phase in Fig. 2. Generalizing, along the lines of constant physics the slope of the line of measured values of the pseudoscalar mass is a direct measure of the sign of the beta function. Fig. 6 provides evidence for the Coulomb phase, with a positive sign of the beta function, in full agreement with the more naive discussion of Fig. 5(b).




FIG. 5: (color online) (a) Measurements of the pseudoscalar (blue) and vector (red) masses versus lattice coupling at several values of the bare quark mass, from bottom to top $am = 0.04, 0.05, 0.06$ and 0.07 . Lines displayed represent a global parametrization, with a mixed $O(m)$ polynomial quark mass dependence and $O(\beta^2)$ polynomial dependence, with lattice parameter $\beta = 6/g_L^2$, and producing a reduced χ^2 per d.o.f. just over unity for both channels. Errors include fitting systematics from combining several methods. (b) The measured π to ρ mass ratio as a function of the bare mass and decreasing coupling g_L , bottom to top $6/g_L^2 = 3.5$ to 4 . The superimposed lines are ratios of the best fits in Fig. 5(a).




FIG. 6: Pseudoscalar mass along lines of constant physics. The physical pion mass is identical along lines of constant physics, such that a decreasing pion mass in lattice units for increasing g_L , implies an increase of the lattice spacing for weaker couplings, signature of the Coulomb phase. The pseudoscalar mass along lines of constant physics was constructed by interpolating along the isolines of the ratio of the separate interpolations of pseudoscalar and vector masses. Different polynomial interpolations produced compatible results, and in agreement with a non-interpolated analysis of raw data. Labels give the value of the ratio m_π/m_ρ along the isolines.

To summarize, for an $SU(3)$ gauge theory with twelve flavors the lattice phase transition from a chirally broken to a symmetric phase is of a non-thermal nature, as is expected for a theory within the conformal window. A study of the spectrum in the symmetric phase close to the transition shows a positive sign of the beta function, incompatible with asymptotic freedom and implying the emergence of a Coulomb phase. Assuming the validity of the scenario in Refs. [3, 4] and Fig. 2, the observation of a Coulomb phase points to the existence of an infrared stable fixed point for the theory with twelve flavors and of a conformal window.

This work was in part based on the MILC public lattice gauge theory code. We thank M. Bochicchio, F. Di

Renzo, F. Sannino, C. deTar for comments and discussions. Computer time was provided through the Dutch National Computing Foundation (NCF) and the University of Groningen.

-
- [1] W. E. Caswell, Phys. Rev. Lett. **33**, 244 (1974).
 - [2] T. Banks and A. Zaks, Nuc. Phys. B **196**, 189 (1982).
 - [3] T. Appelquist, J. Terning, and L. Wijewardhana, Phys. Rev. Lett. **77**, 1214 (1996).
 - [4] V. A. Miransky and K. Yamawaki, Phys. Rev. D **55**, 5051 (1997).
 - [5] T. Appelquist, G. T. Fleming, and E. T. Neil, Phys. Rev. Lett. **100**, 171607 (2008).
 - [6] A. Deuzeman, M. P. Lombardo, and E. Pallante, Phys. Lett. B **670**, 41 (2008).
 - [7] G. T. Fleming, PoS(LAT2008)**21**.
 - [8] T. Appelquist, G. T. Fleming, and E. T. Neil, Phys. Rev. D **79**, 076010 (2009).
 - [9] X.-Y. Jin and R. D. Mawhinney, PoS(LAT2008)**59**.
 - [10] Z. Fodor, K. Holland, J. Kuti, D. Nogradi, and C. Schroeder, arXiv:0911.2463.
 - [11] A. Hasenfratz, arXiv:0911.0646.
 - [12] X.-Y. Jin and R. D. Mawhinney, PoS(LAT2009)**49**.
 - [13] T. Appelquist, A. G. Cohen, and M. Schmaltz, Phys. Rev. D **60**, 045003 (1999).
 - [14] J. Braun and H. Gies, J. High Energy Phys. **2006**, 024 (2006).
 - [15] T. A. Ryttov and F. Sannino, Phys. Rev. D **76**, 105004 (2007).
 - [16] F. Sannino, arXiv:0804.0182.
 - [17] P. Damgaard, U. Heller, A. Krasnitz, and P. Olesen, Phys. Lett. B **400**, 169 (1997).
 - [18] D. B. Kaplan, J.-W. Lee, D. T. Son, and M. A. Stephanov, arXiv:0905.4752.
 - [19] A. Kocic, S. Hands, J. B. Kogut, and E. Dagotto, Nucl. Phys. B **347**, 217 (1990).